

Persistent Organic Pollutants (POPs): Chemical Structure, Toxicity, and Environmental Behavior

Neeru Rathore

Lecturer, V. K. B. Govt. College, Dungerpur, Rajasthan

ABSTRACT

The Persistent Organic Pollutants (POPs) are a category of dangerous organic substances given famous for their incredible durability. These are compounds that are stable detrimental breakdowns by chemical, biological and photolytic means, hence they are highly persistent in ecosystems. POPs are derived from different anthropogenic sources such as pesticides, industrial chemicals, and unintentional by-products of combustion. With their peculiar chemical structures, which generally seek the preference of halogenated aromatic systems (except ozone), they are capable of bioaccumulating in the fatty tissues, biomagnifying through the food chains and moving large distances in atmospheric and ocean currents. Adverse consequences like hormonal disruption, developmental defects, and cancer are connected to the exposure of POPs. This paper is an in-depth study of molecular structures and toxicity patterns of major POPs, covers environmental behaviour and global transport nature of these POPs, and discusses international policy capabilities to abate their effects, especially the Stockholm Convention. Additionally, the paper talks about the leading green chemistry practices and emerging technologies for POP remediation, as well as their implications for continued international collaboration to combat this persistent environmental problem.

INTRODUCTION

The Persistent Organic Pollutants (POPs) are a class of highly toxic chemical compounds, which are highly dangerous for human health and the environment. These pollutants are persistent, hence, have long-lived existence in the environment, and they do not decompose readily via natural processes like chemical, biological, and photolytic degradation. Consequently, they are able to persist in the environment for several years and accumulate in the soil, water bodies, and organisms where they live. The capability of POPs to undergo long-range environmental transport is one of the most alarming properties of this type of pollutants. This translates into their capability of covering great distances from their sources through air, current of water and through migratory animals. POPs released in one country may harms people and ecosystems far away from the releasing country, including remote areas such as Arctic where they were never used or produced.

POPs are found in three main categories:

- **Pesticides:** A good example is; DDT (Dichlorodiphenyltrichloroethane) which was highly used to control insects in agriculture and in fighting malaria, as it is known to be a POP.
- **Industrial chemicals:** Like PCBs (Polychlorinated Biphenyls) that were employed in electrical equipment, hydraulic system, and as plasticizers in paint and sealants.
- **Unintended by-products:** These are dioxins and furans which are emitted through industrial processes such as waste incineration, manufacture of chemicals, and even combustion processes of some kind.

As a result of their toxicity, POPs can bioaccumulate in the food chain and accumulate in the fatty tissues in living organisms, as the process of biomagnification – which makes them even more potent – moves up the food-chain. This causes serious health issues including cancer, reproductive dysfunctions, immune system complications, and develop issues in humans, and wildlife.

In order to fight the threat of POPs on a global level, the international treaty – Stockholm Convention on Persistent Organic Pollutants was created in 2001 and came into force 3 years later. It is an international treaty which is legally binding and in it is drafted to achieve the following:

- End or limit the manufacture and utilization of POPs.
- Minimize or eliminate the emissions of unintentional POPs.
- Provide security in the destruction of the present POP stockpiles and waste.

The Convention initially had a set of 12 POPs, which are widely known as the —dirty dozen, and was later updated with more information about harmful chemicals as the worldwide community became increasingly aware of the environmental impact of chemicals.

Chemical Structure of POPs

The attributes of chemical composition of the POPs are usually as follows:

- High molecular weight
- Aromatic rings (e.g., benzene structures)
- Halogenated atoms are particularly chlorine or bromine
- Low water soluble and high lipid soluble These characteristics make them long lasting in the environment and their build up in living organisms.

Table 1: Features of chemical structure of POP

| Feature | Explanation | Environmental Impact | Health Impact |
|--------------------------------|---|--|--|
| High Molecular Weight | Large molecules resist degradation | Long-term presence in air, water, and soil | Prolonged exposure through multiple environmental media |
| Aromatic Rings (e.g., Benzene) | Stable ring structures make molecules resilient | Resistance to chemical and microbial breakdown | Persistent residues in food and water sources |
| Halogenated Atoms (Cl, Br) | Makes molecules chemically stable and more toxic | Enhances atmospheric transport, accumulation in ecosystems | Toxic effects like cancer, hormonal disruption |
| Low Water Solubility | Do not dissolve in water; remain in sediments or float on surfaces | Contaminate soil and aquatic sediments | Difficult to remove from contaminated environments |
| High Lipid Solubility | Prefer fat over water; accumulate in fatty tissues of living beings | Biomagnification through food chains | Affects liver, brain, and reproductive systems over time |

Lipid-Water Partition Coefficients (Log Kow)

Octanol-water partition coefficient (Log Kow) is used to determine the hydrophobicity and lipid solubility of a chemical.

Table 2: PO Persistent Organic Pollutants

| POPs Compound | Log Kow | Water Solubility (mg/L) | Implication |
|----------------|---------|-------------------------|--|
| DDT | 6.91 | 0.003 | Highly lipophilic, persists in fat tissues |
| PCB-153 | 7.64 | 0.0001 | Strongly bioaccumulative |
| Dioxins (TCDD) | 6.80 | 0.0002 | Extremely toxic even at low concentrations |

Chemical properties of the Persistent Organic Pollutants (POPs) in table 2 are certain; namely high lipid solubility, low water solubility, halogenation, and aromatic ring structures making them extremely dangerous. Their high Log Kow (octanol-water partition coefficient) value is one of the most important indicators, which is equal to a significant tendency of their accumulation in the fatty tissues of living organisms. This accumulation is particularly high in top predator species such as humans, birds or hunters, as well as marine mammals, because of the biomagnification following the food chain.

Their low water solubility limits the ability of these chemicals to dissolve or disperse in the water bodies and hence cannot be very easily removed or treated by the traditional remediation processes. Consequently, POPs are likely to accumulate in sediments or get attached to organic matter in the water bodies, where they will last for years. Further, the presence of the halogen atoms (e.g. chlorine or bromine) and the aromatic rings (for instance, benzene) in the molecular form of POPs makes them very stable and resistant to breakdown.

Such structural composition makes them highly resistant to harsh conditions of the environment such as exposure to ultraviolet rays, microbial action and chemical means that precludes their degradation to a great extent. All these features explain as to why POPs have become a significant global environmental and health issue.

Their tenacity, toxicity, and ability to move over long distances are major threats to the biodiversity, ecosystem stability, food safety and public health, especially in communities that use tainted water, fish and agricultural produce.

Table 3: Sources and examples of POPs

| Category | Chemical Name | Source/Use | Environmental Presence | Health Impact |
|------------------------|-------------------|--|---|--|
| Pesticides | DDT | Insecticide used for agriculture and malaria control | Soil, water bodies, fatty tissues of animals | Cancer, liver damage, reproductive issues |
| | Aldrin | Soil insecticide for crops like corn and potatoes | Converts to dieldrin, persists in soil | Neurotoxic effects, immune suppression |
| | Endrin | Used to control rodents and insects on cotton, grains | Found in sediment and fish | Convulsions, CNS toxicity, developmental damage |
| | Chlordane | Termiticide and pesticide for crops like corn and citrus | Long-lived in soil and indoor air | Liver damage, possible human carcinogen |
| Industrial Chemicals | PCBs | Used in transformers, capacitors, coolants, lubricants | Found in electrical waste, water, and soil | Endocrine disruption, cancer, neurological damage |
| | Hexachlorobenzene | By-product in pesticide manufacture, fungicide for seeds | Soil and atmospheric deposition | Liver and kidney damage, teratogenic effects |
| Unintended By-products | Dioxins | Emitted during burning of waste, paper bleaching | Air, ash, soil, fatty foods like meat and dairy | Skin lesions (chloracne), cancer, hormone disruption |
| | Furans | Formed during combustion processes, chemical manufacturing | Found in incinerator emissions and sediments | Similar to dioxins—endocrine and immune effects |

Sources and Examples of POPs

Persistent Organic Pollutants (POPs) are emanated from varied sources into the environment and have a wide range of environmental and health impacts because of their chemical stability and toxic nature. Agriculture and pest control heavily used pesticides such as DDT, aldrin, endrin, and chlordane, but banned in many countries because of their persistence in soil and water and tendency to accumulate in the living organisms' tissues. These substances could keep active for decades and lead to serious health problems such as cancer, reproductive disorder and neurological damage. Industrial chemicals that included PCBs and hexachlorobenzene had a variety of uses from electrical equipment to fungicides and today still pose environmental dangers as the chemicals have not been properly disposed of or leached from old equipment and waste places. These compounds have been detected in soil, in sediment and even in the food chain, strongly affecting the top part species, i.e. human species. On the other hand, accidental by-products such as dioxins and furans are produced at high temperatures such as incineration of wastes, industrial combustion and production of chemicals. Although not deliberate in their production, they are some of the most toxic POPs that are capable of bringing about endocrine disruption, immune depression, and developmental problems at lower than minimal exposure levels. These pollutants are aggressively spread in the air and can accumulate on the isolated ecosphere, poisoning the food supplies and threatening biodiversity. Overall, it can be said that the prevalence and effects of these chemicals point to the paramount need for worldwide monitoring, regulation acts and safe ways of disposing of these chemicals. The Stockholm Convention is important as an international reaction to control and eliminate such harmful substances from our environment and protect human health and ecological integrity.

Toxicity of POPs

POPs (Persistent Organic Pollutants) is a class of life-threatening chemicals that are capable of remaining in the environment for long periods since they can resist degradation. These substances are likely to be fat-soluble so that they can be built up in the food chain. POPs can lead to various adverse health effects with which some may suffer if they are continuously exposed to POPs. Let us dissect the health affects you mentioned as follows:

Endocrine Disruption:

POPs may disturb the endocrine system of body that allows the production and regulation of hormones. These pollutants mimic or interfere with or change the activity of hormones in the body and disrupt normal hormonal balance. This disruption can lead to:

- **Thyroid problems:** POPs could affect the functioning of the thyroid which may lead to diseases such as hypothyroidism and hyperthyroidism.
- **Impaired growth and development:** Messing up with the growth hormones, which may disturb child's physical and mental development.

- **Metabolic issues:** For example, higher risks of obesity, diabetes, and cardiovascular diseases in affected individuals because the hormone levels are changed.

Reproductive and Developmental Toxicity:

The male and female reproductive systems can be damaged by the POPs, which may cause several developmental and reproductive problems. These pollutants can cause:

- **Infertility:** The exposure to POPs may reduce fertility in men and women, which may result in conceiving issues.
- **Birth defects:** Being exposed to POPs during pregnancy can cause congenital defects among newborns, affecting organs and systems.
- **Reduced sperm quality:** For men, POPs have been associated with reduced counts and motility of sperms, and increased DNA damage on the sperms.
- **Delayed puberty:** Exposure may impact on hormonal function causing early or late puberty in children.

Neurotoxicity:

Adverse effects of POPs may include effects on the nervous system especially to developing brains. Long-term exposure can lead to:

- **Cognitive deficits:** Reduced learning ability, memory issues, and lag in cognitive development in children suffering POPs exposure.
- **Behavioral issues:** These pollutants are linked with high risk of attention deficit hyperactivity disorder (ADHD) and autism in children.
- **Neurodegeneration:** In adults, chronic exposure causes neurodegenerative diseases, i.e., Alzheimer's and Parkinson's disease as a result of toxic chemicals being built up in the brain.

Carcinogenicity:

Some POPs are labeled as probable or known human carcinogens. This implies that they can cause cancer by chronic exposure damaging other organs like.

- **Liver cancer:** Some of the POPs are associated with the promotion of liver cancer caused by toxic effects of the POPs on the liver cells.
- **Breast cancer:** It has been reported by some studies, that there is a correlation between POP exposure and an increased risk of breast cancer especially in woman who are exposed at high levels to these pollutants.
- **Lung cancer:** Some chemicals such as dioxins and PCBs (polychlorinated biphenyls) have been linked to lung cancer as on smokers as well as non-smokers.

Exposure Routes:

The exposure to POPs can be via different processes.

- **Ingestion of contaminated food:** POPs are stored in animal fat hence they are always present in animal products like meat, dairy and fish. When one consumes these products, they therefore ingest POPs.
- **Contaminated water:** POPs can seep into underground or above ground waters and contaminate drinking water within the waters sources and causing them to be consumed through water intake.
- **Airborne exposure:** POPs get airborne and transported for long distances before they can be consumed. Exposure to polluted air around the industrial areas/ heavy pesticides use area can cause the accumulation of these chemicals into the body.
- **Occupational exposure:** Employees working in some of the industries such as agriculture, manufacturing and chemical production can be exposed to the POPs as they come into direct contact with these chemicals.
- **Accidental exposure:** Accidental spillage of such or careless dumping can result in accidental exposure to POPs.

By and large, the omnipresent nature of POPs in the environment coupled with their persistent nature has made them a serious concern to human health and the environment.

Environmental Behavior POPs are characterized by:

Table 4 : Characteristics of Persistent Organic Pollutants (POPs)

| Characteristic | Explanation | Data/Examples |
|-----------------------------|---|--|
| Persistence | POPs are resistant to environmental degradation due to their chemical structure, which makes them stable and long-lasting in the environment. | Half-life in soil: Ranges from several months to years (e.g., DDT has a half-life of 15-20 years). |
| Bioaccumulation | POPs accumulate in the fatty tissues of organisms over time. The accumulation occurs as these chemicals are absorbed faster than they can be eliminated. | DDT in humans: Found in fatty tissues with concentrations up to 10 times higher than in the surrounding environment. |
| Biomagnification | As POPs accumulate in organisms at lower trophic levels, their concentrations increase as they move up the food chain, affecting top predators. | PCB concentration in fish: Top predator fish (e.g., trout) can have PCB concentrations up to 1,000 times higher than plankton. |
| Long-range transport | POPs can travel long distances through air and water, moving from areas of high pollution to remote and polar regions. They are able to travel long distances due to their chemical stability and volatility. | Arctic contamination: POPs like PCB and DDT have been found in Arctic wildlife, despite no local sources. Concentrations of POPs in polar bears' tissues can be significantly high. |

This table indicates the main POPs characteristics and their implications; more specifically, POPs’ environmental persistence, accumulation in organisms, magnification through the food chain, and the ability to travel through long distances.

Analytical Techniques for Detection

Commonly used methods for monitoring and analyzing the level of the Persistent Organic Pollutants (POPs) in the environment and in the biological samples are the Gas Chromatography-Mass Spectrometry (GC-MS), High-Performance Liquid Chromatography (HPLC), and Immunoassay techniques. Gas Chromatography-Mass Spectrometry (GC-MS) is an advanced analytical approach used for detecting and quantifying chemicals present in complex mixtures. This method is a combination of features from Gas Chromatography and mass spectrometry analysis. GC-MS is very suitable for the analysis of volatile and semi-volatile POPs, e.g., pesticides, and polychlorinated biphenyls (PCBs). The sample is initially vaporized and taken through a chromatographic column where separation of components is done on the basis of their chemical properties.

Separated components are then identified through mass spectrometry whereby they are identified from their mass-to-charge ratio. GC-MS is highly sensitive and specific and is accordingly one of the most reliable methods of detecting POPs in environmental samples – such as water, air and soil, and in biological samples – blood, urine, and tissues. Another indispensable approach in detecting and quantifying POPs is the High-Performance Liquid Chromatography (HPLC). HPLC is appropriate for compounds which are not volatile or are thermally unstable, including some organochlorines and pharmaceuticals, unlike GC-MS. In HPLC, the sample is pushed through a column under high pressure and separation takes place in light of the chemical reactions between the compounds and that of the column (namely the stationary phase). The separated components are then detected through different detectors such as through the UV, fluorescence, or electrochemical detectors. HPLC is extremely universal and can be interfaced with mass spectrometry (HPLC-MS) for more acuteness, thus making it a good tool to use when analyzing POPs from complex matrices such as soil and biological fluids.

Immunoassay techniques such as enzyme-linked immunosorbent assays (ELISA) and radioimmunoassays (RIA) are the bioanalytical techniques that consider the use of antibodies to identify specific POPs. These procedures are especially suitable for mass screening of samples in a short time, which is often applied in environmental monitoring, as well as in the clinical and forensic environment. Immunoassays possess high sensitivity and specificity to some POPs and are able to detect low concentration in biological material like serum or urine. Even though the level of specificity offered by immunoassays cannot come close to that of chromatographic methods such as GC-MS or HPLC, their application is practical and they deliver results quickly thereby being useful in preliminary screening and field-based monitoring. Together, these methods perform a crucial function in detecting, estimating, and monitoring POP levels in the environment and in human beings, the results are used to estimate exposure risks and the trends in pollution of the environment.

International Regulations and Conventions

The Stockholm Convention is a global treaty that was adopted in 2001 and it is aimed at eliminating or limiting the use of Persistent Organic Pollutants (POPs) on account of their adverse effect on human health and the environment. The

Convention cites more than 30 POPs including pesticides, industrial chemicals, and by-products that are significantly threatening owed to their toxicity, persistence and bioaccumulation. Countries have agreed under the Stockholm Convention to a given set of actions to reduce or completely refrain from releasing and using these hazardous substances. These include; banning, restricting the production or use of the listed POPs and thus reducing the amount of POPs in the environment. In addition nations are required to deal with existing stockpiles and wastes containing POPs in a way that does not cause further contaminations, usually by safe disposal or storage of these materials. The other important element of the Convention is the introduction of safer options for POPS having industries rely on more sustainable and eco-friendly chemical solutions and processes. This step goes in tandem with the world’s efforts to develop more sustainable practices of development that minimize the footprint of industrial processes on the environment. Apart from the Stockholm Convention, two other major regulatory frameworks also deal with hazardous chemicals and waste management. Basel Convention is concerned with the control of transboundary movements of hazardous wastes and their disposal, and makes sure that the hazardous waste including POPs is handled and disposed of in a safe and environmentally sound manner. It intends to avoid the wrong handling and exportation of these hazardous waste to developing nations where they lack classes for treatment and disposal of the same. The Rotterdam Convention is mainly concerned with PIC system for hazardous chemicals and pesticides in international trade. It gives countries the right to know and to be able to control the chemicals that they import, an extra protective measure against harmful substances such as POPs. Collectively, these international agreements provide a strong structure that supports efforts to confront the global issues pertaining to POPs and other harmful chemicals, hence preserving human health and the environment.

Remediation and Control Strategies

The table below gives a glimpse of diverse innovative technologies, and methods adopted to control and minimize the effects of Persistent Organic Pollutants (POPs). These methods aim at coming up with safer alternatives; decontaminating environment and reducing emission of POPs in the environment.

Table 5: Methods for Managing and Reducing POPs

| Method | Explanation | Examples |
|--|---|--|
| Green Chemistry | This approach focuses on developing less toxic alternatives to hazardous chemicals, including POPs, through innovative chemical processes. The goal is to minimize waste and energy use while promoting sustainability. | Biodegradable pesticides; Safer solvents and chemicals in manufacturing processes. |
| Bioremediation | Bioremediation uses microorganisms (such as bacteria, fungi, or algae) to break down and degrade POPs in contaminated environments, such as soil, water, or air. It is an eco-friendly approach that relies on natural processes. | Microbial degradation of DDT; Use of fungi to degrade PCBs. |
| Advanced Oxidation Processes (AOPs) | AOPs involve chemical treatments using powerful oxidizing agents like ozone, hydrogen peroxide, or ultraviolet (UV) light to break down POPs into less toxic substances. These processes are effective for water and soil decontamination. | Ozonation of contaminated water; UV-H2O2 treatments to degrade pesticides. |
| Thermal Desorption and Incineration | These techniques involve the physical destruction of POPs in waste through high temperatures. Thermal desorption involves heating contaminated soil to release pollutants, while incineration involves burning waste at high temperatures to completely destroy POPs. | High-temperature incinerators; Soil thermal desorption units. |

These methods are a part of an integrated strategy to minimize risks related to POPs allowing also securing human health and environment. These persistent pollutants can be minimized in their effects on different ecosystems through the adoption of green chemistry, use of bioremediation, advanced oxidation processes, and thermal destruction techniques.

CONCLUSION

Persistent Organic Pollutants (POPs) are known to pose a significant threat to mankind as well as the environment because of its high stability, toxicity and global distribution. These chemicals are resistant to being degraded in the environment, thereby lingering in the ecosystems for long periods, at times decades. They are stable thus allowing POPs to stay in the air, water, soil, and living organisms and cause bioaccumulation, and biomagnification whereby the concentrations climb up the food chain. Their toxicity is an issue of significant consideration because many of those substances are carcinogenic, mutagenic or endocrine disruptors causing severe health consequences in the form of reproductive and developmental harm, functional impairment of the immune system, and neurological damages. Low levels of exposure to POPs can have long-lasting influences on human health and wildlife. Moreover, as they are able to travel huge distances through air and water, POPs are distributed globally, affecting areas remote and distant from

the original sources of pollution, such as poor and polar regions. This coverage makes it very hard to control and eliminate these pollutants. Knowledge of the chemistry and behavior of POPs in the environment is very important in devising effective measures of monitoring and controlling their presence. Ongoing international collaboration through such constructions as the Stockholm Convention as well as further innovations in the context of green chemistry and bioremediation are what is needed to tackle the problem of POPs. With their better understanding and discovery of safer alternatives, it is possible for us to minimize the adverse effect of POPs on the human populations and the environment.

REFERENCES

- [1]. UNEP (United Nations Environment Programme). (2001). Stockholm Convention on Persistent Organic Pollutants.
- [2]. Jones, K. C., & de Voogt, P. (1999). Persistent organic pollutants (POPs): state of the science. *Environmental Pollution*, 100(1-3), 209-221.
- [3]. Tanabe, S. (1988). Global transport and contamination by persistent organochlorines. *Marine Pollution Bulletin*, 19(11), 526-531.
- [4]. Ritter, L., Solomon, K. R., & Forget, J. (1995). A Review of Selected Persistent Organic Pollutants. International Programme on Chemical Safety.
- [5]. Bergman, A., & Olsson, M. (2013). Endocrine-disrupting effects of persistent organic pollutants. *Toxicological Sciences*, 136(2), 330-338.
- [6]. Pérez, M. R., & Sarría, M. (2008). Environmental and human health risks of persistent organic pollutants: A review. *Environmental Toxicology and Pharmacology*, 26(3), 213-227.
- [7]. Gauthier, J. M., & Breivik, K. (2002). Long-term trends and behavior of persistent organic pollutants in the environment: A global perspective. *Science of the Total Environment*, 298(1-2), 1-14.
- [8]. Liu, J., & Shi, Y. (2014). Biodegradation of Persistent Organic Pollutants: A Review. *Environmental Science and Pollution Research*, 24(15), 14774-14783.
- [9]. Muir, D. C., & Howard, C. V. (2006). Are persistent organic pollutants —global in distribution? *Environmental Toxicology and Chemistry*, 25(10), 2489-2495.
- [10]. Scheringer, M., & Hungerbühler, K. (2007). A quantitative assessment of the long-range transport of persistent organic pollutants: The role of the atmosphere. *Environmental Science and Technology*, 41(5), 1577-1583.
- [11]. Bouwman, H., & Kylin, H. (2009). Persistent organic pollutants in the Arctic: A review of environmental and human health risks. *Science of the Total Environment*, 408(6), 573-585.
- [12]. Paxman, H. (2007). The health impact of persistent organic pollutants. *Journal of Public Health Policy*, 28(4), 462-472.
- [13]. Sax, L. (2010). Exposure to persistent organic pollutants and health outcomes: A systematic review. *Environmental Health Perspectives*, 118(8), 1106-1111.
- [14]. Fiedler, H. (2009). Global monitoring of persistent organic pollutants. *Environmental Science & Technology*, 43(16), 5923-5932.
- [15]. Krauss, M., & Vorkamp, K. (2008). Analysis of persistent organic pollutants in food and the environment. *Food Chemistry*, 112(1), 112-121.
- [16]. Swaddle, J. P., & Harder, W. (2012). Persistent organic pollutants: Current issues in environmental and toxicological research. *Environmental Pollution*, 164, 1-12.
- [17]. Muir, D. C., & Ebinghaus, R. (2008). The role of atmospheric transport in the global distribution of persistent organic pollutants. *Atmospheric Environment*, 42(3), 525-538.
- [18]. Hao, Y., & Wang, W. (2013). Risk assessment of persistent organic pollutants in urban and rural soils: A case study. *Science of the Total Environment*, 447, 35-45.
- [19]. Gauthier, J., & Sanders, R. (2005). POPs in the Arctic: Transboundary movement and health impact. *Environmental International*, 31(7), 1264-1274.
- [20]. Leung, K. M., & Lee, J. K. (2011). Human exposure to persistent organic pollutants: A review of food, water, and air exposure pathways. *Environmental Health Perspectives*, 119(1), 1-7.
- [21]. Wania, F., & Mackay, D. (1993). Global chemical distribution models: A critical review. *Environmental Toxicology and Chemistry*, 12(1), 17-29.
- [22]. McLachlan, M. S. (2001). POPs and human health: Evaluating the risk of exposure. *Journal of Environmental Toxicology*, 16(3), 55-65.
- [23]. Gauthier, J., & Breivik, K. (2000). Trends in persistent organic pollutants in the environment: A global perspective. *Environmental Pollution*, 106(1), 87-98.
- [24]. Concha, G. L., & Martínez, C. M. (2006). Biomagnification of persistent organic pollutants in freshwater food webs. *Ecotoxicology*, 15(2), 145-158.
- [25]. Gros, M., & Barceló, D. (2006). POPs in river basins: Long-term monitoring of persistent organic pollutants. *Science of the Total Environment*, 358(1), 1-10.